European Journal of Pharmacology, 236 (1993) 503–506
© 1993 Elsevier Science Publishers B.V. All rights reserved 0014-2999/93/$06.00

Author index

Abacioğlu, N., see Çakici, Í. (236) 97
Abbenante, J., see Kerr, D.I.B. (236) 239
Abe, Y., see Tamaki, T. (236) 15
Abiko, Y., see Hoque, A.N.E. (236) 269
Ahlenius, S., see Salmi, P. (236) 395
Ahrén, B., see Skoglund, G. (236) 69
Aki, Y., see Tamaki, T. (236) 15
Allard, B. and Lazdunski, M.
Pharmacological properties of ATP-sensitive K^+ channels in mammalian skeletal muscle cells (236) 419
Allerup, P., see Glenthøj, B. (236) 401
Amorim, C.Z., Cordeiro, R.S.B. and Vargaftig, B.B.
Interference of PCA 4248, a novel PAF receptor antagonist, with antigen-induced paw edema in mice (236) 301
Arányi, P., see Horváth, E.J. (236) 151
Armario, A., see Martí, J. (236) 295

Bai, X., see Walker, J.M. (236) 495
Baptista, T., Weiss, S.R.B. and Post, R.M.
Carbamazepine attenuates cocaine-induced increases in dopamine in the nucleus accumbens: an in vivo dialysis study (236) 39
Barber, A.
μ- and κ-opioid receptor agonists produce peripheral inhibition of neurogenic plasma extravasation in rat skin (236) 113
Barry, D.I., see Glenthøj, B. (236) 401
Beer, M.S., Stanton, J.A., Hawkins, L.M. and Middlemiss, D.N.
5-Carboxamidotryptamine-insensitive 5-HT_1-like receptors are concentrated in guinea pig but not rat, claustrum (236) 167
Berkenkopf, J.W., see Grimes, D. (236) 217
Bernard, J.-F., see Huang, G.-F. (236) 449
Besson, J.-M., see Huang, G.-F. (236) 449
Bojö, L., Lefebvre, R.A., Nellgård, P. and Cassuto, J.
Involvement of vasoactive intestinal polypeptide in gastric reflex relaxation (236) 443
Bolwig, T.G., see Glenthøj, B. (236) 401
Bonini, I., see Sacchetti, G. (236) 171
Bonnot-Lours, S., see Tavitian, B. (236) 229
Borsodi, A., see Fürst, Z. (236) 209
Boulton, C.L., McCrohan, C.R. and O'Shaughnessy, C.T.
Cyclic AMP analogues increase excitability and enhance epileptiform activity in rat neocortex in vitro (236) 131
Bouysset, F., see Vayssettes-Courchay, C. (236) 121
Bowen, W.D., see Walker, J.M. (236) 495
Briley, M., see Marien, M. (236) 487
Britton II, L.W., see Weishaar, R.E. (236) 363
Bruhn, T., see Glenthøj, B. (236) 401
Búzás, B., see Fürst, Z. (236) 209

Çakici, Í., Tunçtan, B., Abacioğlu, N. and Kanzik, Í.
Epithelium-dependent responses of serotonin in a co-axial bioassay system (236) 97
Callahan, P.E., see Ratz, P.H. (236) 433
Carlson, R.P., see Grimes, D. (236) 217
Carroll, F.I., see Walker, J.M. (236) 495
Cassuto, J., see Bojö, L. (236) 443
Castle, N.A., Haylett, D.G., Morgan, J.M. and Jenkinson, D.H.
Dequalinium: a potent inhibitor of apamin-sensitive K^+ channels in hepatocytes and of nicotinic responses in skeletal muscle (236) 201
Chang, K.C., Ko, H.J., Cho, S.D., Yoon, Y.J. and Kim, J.H.
Pharmacological characterization of effects of verapamil and GS 283 on isolated guinea pig and rat trachealis (236) 51
Chase, T.N., see Engber, T.M. (236) 385
Cho, S.D., see Chang, K.C. (236) 51
Chuluyan, H.E., see Saphier, D. (236) 183
Code, R.A., see Tang, A.H. (236) 1
Cohen, D.S., see Olson, R.W. (236) 75
Colpaert, F., see Marien, M. (236) 487
Connor, H.E., see O'Shaughnessy, C.T. (236) 319
Cools Waeterloos, G., see Sacchetti, G. (236) 171
Cordeiro, R.S.B., see Amorim, C.Z. (236) 301
Cowan, A., see Wheeler-Aceto, H. (236) 193
Crouzel, C., see Tavitian, B. (236) 229
Cumming-Hood, P.A., Strahlendorf, H.K. and Strahlendorf, J.C.
Effects of serotonin and the 5-$HT_{2/1C}$ receptor agonist DOI on neurons of the cerebellar dentate/interpositus nuclei: possible involvement of a GABAergic interneuron (236) 457

De Costa, B.R., see Hudzik, T.J. (236) 279
Di Giamberardino, L., see Tavitian, B. (236) 229
Doolette, D.J., see Kerr, D.I.B. (236) 239
Dotson, R., see Olson, R.W. (236) 75
Duckles, S.P., see Li, Y. (236) 373

Egan, J.W., see Smith III, E.F. (236) 477
Engber, T.M., Marin, C., Susel, Z. and Chase, T.N.
Differential effects of chronic dopamine D_1 and D_2 receptor agonists on rotational behavior and dopamine receptor binding (236) 385
Engel, J.A., see Heilig, M. (236) 339
Evans, D.L., see Lahti, R.A. (236) 483

Falch, E., see White, H.S. (236) 147
Fekete, M.I.K., see Horváth, E.J. (236) 151
Ferraris, V.A., see Weishaar, R.E. (236) 363
Figur, L.M., see Lahti, R.A. (236) 483
Filep, J.G., see Hermán, F. (236) 143
Francisco, T., see Mendelson, S.D. (236) 177
Franklin, P.H., see Roth, J.E. (236) 327
Franks, R., see Guard, S. (236) 496
Friedmann, T., see Fürst, Z. (236) 209
Fuji, K., Hiramatsu, M., Kameyama, T. and Nabeshima, T.
Effects of repeated administration of propentofylline on memory impairment produced by basal forebrain lesion in rats (236) 411
Fujii, T., see Murai, M. (236) 7
Fukui, T., see Tamaki, T. (236) 15
Fürst, Z., Búzás, B., Friedmann, T., Schmidhammer, H. and Borsodi, A.
Highly potent novel opioid receptor agonist in the 14-alkoxymetopon series (236) 209

Furukawa, S., Satoh, K. and Taira, N.
Opening of ATP-sensitive K^+ channels responsible for adenosine A_2 receptor-mediated vasodepression does not involve a pertussis toxin-sensitive G protein (236) 255

Gagnon, R.C., see Smith III, E.F. (236) 477
García-Sevilla, J.A., see Olmos, G. (236) 467
Garro, M.A., see Olmos, G. (236) 467
Giralt, M.T., see Olmos, G. (236) 467
Glenthøj, B., Hemmingsen, R., Barry, D.I., Allerup, P., Bruhn, T. and Bolwig, T.G.
Electrical kindling of rats treated discontinuously or continuously with haloperidol (236) 401
Grimes, D., Sturm, R.J., Marinari, L.R., Carlson, R.P., Berkenkopf, J.W., Musser, J.H., Kreft, A.F. and Weichman, B.M.
WY-50,295 tromethamine, a novel, orally active 5-lipoxygenase inhibitor: biochemical characterization and antiallergic activity (236) 217
Griswold, D.E., see Smith III, E.F. (236) 477
Gross, R., see Skoglund, G. (236) 69
Guard, S., Pain, D., Franks, R. and Watling, K.J.
[^{125}I]Neurokinin A labels pharmacologically distinct populations of NK_2 binding sites in hamster and rabbit urinary bladder. European Journal of Pharmacology, 232 (1993) 287–290 (236) 496
Gwilt, M., Norton, B. and Henderson, C.G.
Pharmacological studies of K^+ loss from ischaemic myocardium in vitro: roles of ATP-dependent K^+ channels and lactate-coupled efflux (236) 107

Hagiwara, D., see Murai, M. (236) 7
Hamra, M. and Rosen, M.R.
The influence of pH on the use-dependent effects of lidocaine in adult and neonatal canine Purkinje fibers. European Journal of Pharmacology, 230 (1993) 167–175 (236) 497
Hashimoto, K. and London, E.D.
Further characterization of [^{3}H]ifenprodil binding to σ receptors in rat brain (236) 159
Hawkins, L.M., see Beer, M.S. (236) 167
Haylett, D.G., see Castle, N.A. (236) 201
Hayward, N.J., McKnight, A.T. and Woodruff, G.N.
Brain temperature and the neuroprotective action of enadoline and dizocilpine in the gerbil model of global ischaemia (236) 247
He, H., see Tamaki, T. (236) 15
Heath, A.R., see Potter, D.E. (236) 61
Heilig, M., Engel, J.A. and Söderpalm, B.
C-fos antisense in the nucleus accumbens blocks the locomotor stimulant action of cocaine (236) 339
Hemmingsen, R., see Glenthøj, B. (236) 401
Henderson, C.G., see Gwilt, M. (236) 107
Hermán, F., Yano, M. and Filep, J.G.
The in vivo antiaggregatory action of endothelin-1 is not mediated through the endothelin ET_A receptor (236) 143
Hertting, G., see Lauth, D. (236) 165
Hibbs, M.J., see Smith III, E.F. (236) 477
Hillegass, L.M., see Smith III, E.F. (236) 477
Himes, C.S., see Tang, A.H. (236) 1
Hiramatsu, M., see Fuji, K. (236) 411
Ho, I.K., see Tseng, Y.T. (236) 23
Hofman, G., see Van Oosterhout, A.J.M. (236) 379
Hoque, A.N.E., Nasa, Y. and Abiko, Y.
Cardioprotective effect of d-propranolol in ischemic-reperfused isolated rat hearts (236) 269
Horváth, E.J., Hudák, J., Palkovits, M., Lenkei, Z., Fekete, M.I.K. and Arányi, P.
A novel specific binding site for homophthalazines in the rat brain (236) 151
Huang, G.-F., Besson, J.-M. and Bernard, J.-F.
Intravenous morphine depresses the transmission of noxious messages to the nucleus centralis of the amygdala (236) 449
Hudák, J., see Horváth, E.J. (236) 151
Hudzik, T.J., De Costa, B.R. and McMillan, D.E.
σ Receptor-mediated emetic response in pigeons: agonists, antagonists and modifiers (236) 279
Hunt, J., see White, H.S. (236) 147

Iizuka, Y., see Yamaguchi, T. (236) 43
Ikari, N., see Murai, M. (236) 7
Ikeda, T., see Yoshioka, M. (236) 491
Itoh, J., Ukai, M. and Kameyama, T.
Dynorphin A-(1–13) markedly improves scopolamine-induced impairment of spontaneous alternation performance in mice (236) 341
Itoh, K., see Yamaguchi, T. (236) 43
Itoh, N., see Yayama, K. (236) 337
Iwao, H., see Tamaki, T. (236) 15

Jackisch, R., see Lauth, D. (236) 165
Jenkinson, D.H., see Castle, N.A. (236) 201
Jimenez, P., see Salmi, P. (236) 395
Jobert, A., see Tavitian, B. (236) 229

Kadowitz, P.J., see Santiago, J.A. (236) 315
Kameyama, T., see Fuji, K. (236) 411
Kameyama, T., see Itoh, J. (236) 341
Kanzik, İ., see Çakici, İ. (236) 97
Kashimoto, T., see Tachikawa, E. (236) 355
Kawao, M., see Yayama, K. (236) 337
Kerr, D.I.B., Ong, J., Doolette, D.J., Abbenante, J. and Prager, R.H.
3-Amino-2-(4-chlorophenyl)-nitropropane is a new $GABA_B$ receptor agonist, more active peripherally (236) 239
Kim, J.H., see Chang, K.C. (236) 51
Kimura, S., see Tamaki, T. (236) 15
Kirker, M.L., see Weishaar, R.E. (236) 363
Ko, H.J., see Chang, K.C. (236) 51
Kreft, A.F., see Grimes, D. (236) 217
Krogsgaard-Larsen, P., see White, H.S. (236) 147
Kuroume, T., see Yokoyama, T. (236) 427

Lahti, R.A., Evans, D.L., Stratman, N.C. and Figur, L.M.
Dopamine D_4 versus D_2 receptor selectivity of dopamine receptor antagonists: possible therapeutic implications (236) 483
Lattanzio, Jr., F.A., see Ratz, P.H. (236) 433
Laubie, M., see Vayssettes-Courchay, C. (236) 121
Lauth, D., Hertting, G. and Jackisch, R.
Involvement of nitric oxide synthase in 3,4-diaminopyridine-evoked noradrenaline release in rat hippocampus (236) 165
Lazdunski, M., see Allard, B. (236) 419
Lefebvre, R.A., see Bojö, L. (236) 443
Lenkei, Z., see Horváth, E.J. (236) 151
Li, Y. and Duckles, S.P.
Effect of age on vascular content of calcitonin gene-related peptide and mesenteric vasodilator nerve activity in the rat (236) 373
London, E.D., see Hashimoto, K. (236) 159
Löscher, W., see Richter, A. (236) 289
Lou, Y.-P., see Satoh, H. (236) 367
Loubatières-Mariani, M.-M., see Skoglund, G. (236) 69
Lundberg, J.M., see Satoh, H. (236) 367

Maeda, Y., see Murai, M. (236) 7
Maggi, C.A., see Patacchini, R. (236) 31

March, D. and Shaw, C.A.
Fetal, neonatal and adult expression of benzodiazepine receptor subtypes in human visual cortex (236) 333
Marien, M., Briley, M. and Colpaert, F.
Noradrenaline depletion exacerbates MPTP-induced striatal dopamine loss in mice (236) 487
Marin, C., see Engber, T.M. (236) 385
Marinari, L.R., see Grimes, D. (236) 217
Martí, J. and Armario, A.
Effects of diazepam and desipramine in the forced swimming test: influence of previous experience with the situation (236) 295
Mascarella, S.W., see Walker, J.M. (236) 495
Mathis, J., see Olson, R.W. (236) 75
Matsuda, Y., see Nozawa, M. (236) 263
Matsuo, M., see Murai, M. (236) 7
McCrohan, C.R., see Boulton, C.L. (236) 131
McKnight, A.T., see Hayward, N.J. (236) 247
McMillan, D.E., see Hudzik, T.J. (236) 279
Menargues, A., see Olmos, G. (236) 467
Mendelson, S.D., Quartermain, D., Francisco, T. and Shemer, A.
5-HT_{1A} receptor agonists induce anterograde amnesia in mice through a postsynaptic mechanism (236) 177
Middlemiss, D.N., see Beer, M.S. (236) 167
Miralles, A., see Olmos, G. (236) 467
Miyake, H., see Murai, M. (236) 7
Miyaoka, T., see Tseng, Y.T. (236) 23
Miyatake, A., see Tamaki, T. (236) 15
Mizuma, K., see Tachikawa, E. (236) 355
Morgan, J.M., see Castle, N.A. (236) 201
Morikawa, A., see Yokoyama, T. (236) 427
Mukaiyama, O., see Yamaguchi, T. (236) 43
Murai, M., Maeda, Y., Hagiwara, D., Miyake, H., Ikari, N., Matsuo, M. and Fujii, T.
Effects of an NK_1 receptor antagonist, FK888, on constriction and plasma extravasation induced in guinea pig airway by neurokinins and capsaicin (236) 7
Murray, T.F., see Roth, J.E. (236) 327
Musser, J.H., see Grimes, D. (236) 217

Nabeshima, T., see Fuji, K. (236) 411
Nasa, Y., see Hoque, A.N.E. (236) 269
Nellgård, P., see Bojö, L. (236) 443
Nijkamp, F.P., see Van Oosterhout, A.J.M. (236) 379
Nishiyama, A., see Tamaki, T. (236) 15
Nordholm, L., see Sheardown, M.J. (236) 347
Norton, B., see Gwilt, M. (236) 107
Nozawa, M., Toki, S. and Matsuda, Y.
The neuroprotective properties of ES-242s, novel NMDA receptor antagonists, in neuronal cell culture toxicity studies (236) 263

Obach, R., see Olmos, G. (236) 467
Ogidigben, M.J., see Potter, D.E. (236) 61
Okamoto, H., see Yayama, K. (236) 337
Olmos, G., Miralles, A., García-Sevilla, J.A., Giralt, M.T., Pineda, J., Garro, M.A., Ugedo, L., Menargues, A. and Obach, R.
Acute and chronic effects of cholinesterase inhibitors and pilocarpine on the density and sensitivity of central and peripheral α_2-adrenoceptors (236) 467
Olson, R.W., Dotson, R., Mathis, J., Cohen, D.S. and Webb, R.L.
Beneficial effects of combined thromboxane synthase inhibition/receptor blockade with CGS 22652 in a canine model of coronary thrombosis (236) 75
Ong, J., see Kerr, D.I.B. (236) 239
Osei, S.Y., see Santiago, J.A. (236) 315
O'Shaughnessy, C.T. and Connor, H.E.
Neurokinin NK_1 receptors mediate plasma protein extravasation in guinea-pig dura (236) 319
O'Shaughnessy, C.T., see Boulton, C.L. (236) 131

Pain, D., see Guard, S. (236) 496
Palkovits, M., see Horváth, E.J. (236) 151
Pappata, S., see Tavitian, B. (236) 229
Patacchini, R. and Maggi, C.A.
Amastatin interferes with the antagonist properties of MEN 10,208 in the rabbit pulmonary artery but not in the hamster trachea (236) 31
Patrick, S.L., see Walker, J.M. (236) 495
Pearson, H.A. and Wray, D.
Non-competitive antagonism of calcium by magnesium ions at the K^+-depolarised mouse neuromuscular junction (236) 323
Pineda, J., see Olmos, G. (236) 467
Post, R.M., see Baptista, T. (236) 39
Potter, D.E., Ogidigben, M.J. and Heath, A.R.
Ocular actions of an octahydrobenzo[f]quinoline: Ha117 (236) 61
Prager, R.H., see Kerr, D.I.B. (236) 239
Prenant, C., see Tavitian, B. (236) 229

Quartermain, D., see Mendelson, S.D. (236) 177

Ratz, P.H., Callahan, P.E. and Lattanzio, Jr., F.A.
Ketamine relaxes rabbit femoral arteries by reducing $[Ca^{2+}]_i$ and phospholipase C activity (236) 433
Reis, D.J., see Sun, M.-K. (236) 305
Richter, A. and Löscher, W.
The κ-opioid receptor agonist, U50,488H, exerts antidystonic activity in a mutant hamster model of generalized dystonia (236) 289
Rosen, M.R., see Hamra, M. (236) 497
Roth, J.E., Franklin, P.H. and Murray, T.F.
The σ receptor ligand 1,3-di(2-tolyl)guanidine is anticonvulsant in the rat prepiriform cortex (236) 327

Sacchetti, G., Bonini, I., Cools Waeterloos, G. and Samanin, R.
Tianeptine raises dopamine and blocks stress-induced noradrenaline release in the rat frontal cortex (236) 171
Saito, H., see Yoshioka, M. (236) 491
Salmi, P., Jimenez, P. and Ahlenius, S.
Evidence for specific involvement of dopamine D_1 and D_2 receptors in the regulation of body temperature in the rat (236) 395
Samanin, R., see Sacchetti, G. (236) 171
Santiago, J.A., Osei, S.Y. and Kadowitz, P.J.
Inhibitory effects of Hoe 140 on vasodilator responses to bradykinin in the mesenteric vascular bed of the cat (236) 315
Saphier, D., Welch, J.E. and Chuluyan, H.E.
α-Interferon inhibits adrenocortical secretion via μ_1-opioid receptors in the rat (236) 183
Satoh, H., Lou, Y.-P. and Lundberg, J.M.
Inhibitory effects of capsazepine and SR 48968 on citric acid-induced bronchoconstriction in guinea-pigs (236) 367
Satoh, K., see Furukawa, S. (236) 255
Satoh, Y., see Yamaguchi, T. (236) 43
Savelkoul, H.F.J., see Van Oosterhout, A.J.M. (236) 379
Schmidhammer, H., see Fürst, Z. (236) 209
Schott, C.A., Vetrovsky, P. and Stoclet, J.-C.
Cationic amino acids inhibit the effects of L-arginine in rat aorta exposed to lipopolysaccharide (236) 155
Schousboe, A., see White, H.S. (236) 147
Shaw, C.A., see March, D. (236) 333
Sheardown, M.J., Suzdak, P.D. and Nordholm, L.
AMPA, but not NMDA, receptor antagonism is neuroprotective in gerbil global ischaemia, even when delayed 24 h (236) 347
Shemer, A., see Mendelson, S.D. (236) 177
Sim, M.F., see Weishaar, R.E. (236) 363

Skoglund, G., Gross, R., Ahrén, B. and Loubatières-Mariani, M.-M.
Different mechanisms are involved in neuropeptide Y-induced pancreatic vasoconstriction and inhibition of insulin secretion (236) 69
Smith, R.A.G., see Smith III, E.F. (236) 477
Smith III, E.F., Griswold, D.E., Egan, J.W., Hillegass, L.M., Smith, R.A.G., Hibbs, M.J. and Gagnon, R.C.
Reduction of myocardial reperfusion injury with human soluble complement receptor type 1 (BRL 55730) (236) 477
Söderpalm, B., see Heilig, M. (236) 339
Stanton, J.A., see Beer, M.S. (236) 167
Stoclet, J.-C., see Schott, C.A. (236) 155
Strahlendorf, H.K., see Cumming-Hood, P.A. (236) 457
Strahlendorf, J.C., see Cumming-Hood, P.A. (236) 457
Stratman, N.C., see Lahti, R.A. (236) 483
Sturm, R.J., see Grimes, D. (236) 217
Sun, M.-K. and Reis, D.J.
Extracellular H^+ iontophoresis modifies responses to γ-aminobutyric acid and cyanide of reticulospinal vasomotor neurons in rats (236) 305
Susel, Z., see Engber, T.M. (236) 385
Suzdak, P.D., see Sheardown, M.J. (236) 347
Swinyard, E.A., see White, H.S. (236) 147

Tachikawa, E., Yoshinari, H., Takahashi, S., Kashimoto, T., Mizuma, K. and Takahashi, E.
Cisplatin, an antineoplastic drug, inhibits catecholamine secretion from bovine adrenal chromaffin cells (236) 355
Taira, N., see Furukawa, S. (236) 255
Takahashi, E., see Tachikawa, E. (236) 355
Takahashi, S., see Tachikawa, E. (236) 355
Tamaki, T., Nishiyama, A., Yoshida, H., He, H., Fukui, T., Yamamoto, A., Aki, Y., Kimura, S., Iwao, H., Miyatake, A. and Abe, Y.
Effects of EXP3174, a non-peptide angiotensin II receptor antagonist, on renal hemodynamics and renal function in dogs (236) 15
Tang, A.H., Code, R.A. and Himes, C.S.
Antagonism of hypothermia produced by benzodiazepine-related compounds by U-78875 in mice (236) 1
Tavitian, B., Pappata, S., Bonnot-Lours, S., Prenant, C., Jobert, A., Crouzel, C. and Di Giamberardino, L.
Positron emission tomography study of [^{11}C]methyl-tetrahydroaminoacridine (methyl-tacrine) in baboon brain (236) 229
Terada, A., see Yamaguchi, T. (236) 43
Tiseo, P.J. and Yaksh, T.L.
Dose-dependent antagonism of spinal opioid receptor agonists by naloxone and naltrindole: additional evidence for δ-opioid receptor subtypes in the rat (236) 89
Togashi, H., see Yoshioka, M. (236) 491
Toki, S., see Nozawa, M. (236) 263
Tokuyama, K., see Yokoyama, T. (236) 427
Tseng, L.F., see Xu, J.Y. (236) 137
Tseng, Y.T., Miyaoka, T. and Ho, I.K.
Region-specific changes of $GABA_A$ receptors by tolerance to and dependence upon pentobarbital (236) 23
Tujii, H., see Yayama, K. (236) 337
Tunçtan, B., see Çakici, İ. (236) 97

Ugedo, L., see Olmos, G. (236) 467
Ukai, M., see Itoh, J. (236) 341

Van Ark, I., see Van Oosterhout, A.J.M. (236) 379
Van Oosterhout, A.J.M., Van Ark, I., Hofman, G., Savelkoul, H.F.J. and Nijkamp, F.P.
Recombinant interleukin-5 induces in vivo airway hyperresponsiveness to histamine in guinea pigs (236) 379
Vargaftig, B.B., see Amorim, C.Z. (236) 301
Vayssettes-Courchay, C., Bouysset, F., Verbeuren, T.J. and Laubie, M.
Role of the lateral tegmental field in the central sympatho-inhibitory effect of 8-hydroxy-2-(di-n-propylamino)tetralin in the cat (236) 121
Verbeuren, T.J., see Vayssettes-Courchay, C. (236) 121
Vetrovsky, P., see Schott, C.A. (236) 155

Walker, J.M., Bowen, W.D., Patrick, S.L., Williams, W.E., Mascarella, S.W., Bai, X. and Carroll, F.I.
A comparison of (−)-deoxybenzomorphans devoid of opiate activity with their dextrorotatory phenolic counterparts suggests role of σ_2 receptors in motor function. European Journal of Pharmacology, 231 (1993) 61–68 (236) 495
Wallace, A.M., see Weishaar, R.E. (236) 363
Watling, K.J., see Guard, S. (236) 496
Webb, R.L., see Olson, R.W. (236) 75
Weichman, B.M., see Grimes, D. (236) 217
Weishaar, R.E., Kirker, M.L., Wallace, A.M., Ferraris, V.A., Britton II, L.W. and Sim, M.F.
Relationship between inotropic activity and phosphodiesterase inhibition for flosequinan and milrinone (236) 363
Weiss, S.R.B., see Baptista, T. (236) 39
Welch, J.E., see Saphier, D. (236) 183
Wheeler-Aceto, H. and Cowan, A.
Naloxone causes apparent antinociception and pronociception simultaneously in the rat paw formalin test (236) 193
White, H.S., Hunt, J., Wolf, H.H., Swinyard, E.A., Falch, E., Krogsgaard-Larsen, P. and Schousboe, A.
Anticonvulsant activity of the γ-aminobutyric acid uptake inhibitor N-4,4-diphenyl-3-butenyl-4,5,6,7-tetrahydroisoxazolo[4,5-c]pyridin-3-ol (236) 147
Williams, W.E., see Walker, J.M. (236) 495
Wolf, H.H., see White, H.S. (236) 147
Woodruff, G.N., see Hayward, N.J. (236) 247
Wray, D., see Pearson, H.A. (236) 323

Xu, J.Y. and Tseng, L.F.
Increase of nitric oxide by L-arginine potentiates β-endorphin- but not μ-, δ- or κ-opioid agonist-induced antinociception in the mouse (236) 137

Yaksh, T.L., see Tiseo, P.J. (236) 89
Yamaguchi, T., Mukaiyama, O., Itoh, K., Satoh, Y., Terada, A. and Iizuka, Y.
Effects of CS-518, a thromboxane synthase inhibitor, on the asthmatic response (236) 43
Yamamoto, A., see Tamaki, T. (236) 15
Yano, M., see Hermán, F. (236) 143
Yayama, K., Kawao, M., Tujii, H., Itoh, N. and Okamoto, H.
Dup 753 prevents the development of puromycin aminonucleoside-induced nephrosis (236) 337
Yokoyama, T., Tokuyama, K., Morikawa, A. and Kuroume, T.
Bradykinin is less potent in causing airway microvascular leakage in immature than in adult guinea-pigs. Role of neutral endopeptidase (236) 427
Yoon, Y.J., see Chang, K.C. (236) 51
Yoshida, H., see Tamaki, T. (236) 15
Yoshinari, H., see Tachikawa, E. (236) 355
Yoshioka, M., Ikeda, T., Togashi, H. and Saito, H.
Effect of 5-hydroxytryptamine on glossopharyngeal afferent nerve activity in anesthetized rats (236) 491